

Combined population PK modelling of cefepime and enmetazobactam to characterize their individual and combined PK from the phase 1 to 3 studies

Jannik Vollmer (1), Juan Quevedo (2), Nathalie Dunkel (2), Anne Santerre Henriksen (3), Matthias Machacek (1)

(1) LYO-X AG, Basel, Switzerland (2) Advanz Pharma UK, London, United Kingdom (3) Maxel Consulting, Jyllinge, Denmark

Background:

Antimicrobial resistance (AMR) is a global health threat. The most frequent mechanism of resistance in Enterobacterales is the expression of Class A extended-spectrum β -lactamases. Cefepime/enmetazobactam (FEP-META) is a new β -lactam– β -lactamase inhibitor combination with activity against multidrug-resistant Enterobacterales. FEP-META is approved in Europe for the treatment of complicated urinary tract infections (cUTI), hospital-acquired pneumonia, including ventilator-associated pneumonia and bacteraemia that occurs in association with, or is suspected to be associated with, any of the infections listed above.

Goal:

The objective of this study was to characterize the FEP and META population pharmacokinetics (PK) from the combined phase 1, 2 and 3 trials, including the intra-patient correlation of the FEP and META PK parameters.

Methods:

Plasma PK samples were available from a total of 663 subjects in three phase 1 (n=132), one phase 2 (n=43) and one phase 3 (n=488) clinical trial. The population included healthy volunteers (n=113), subjects with renal impairment (n=19) and patients with cUTI (n=531).

A compartmental population PK modelling approach was used. Monolix Suite 2019R2 was used for parameter estimation. For model development, a stepwise model building was employed (**Figure 1**). In the combined population PK model, the covariance of FEP and META random effects of central volumes of distribution and clearances were assessed. To best characterize the between subject variability in the cUTI patient population independent inter-individual variabilities for healthy and cUTI patients were estimated.

Figure 1: Step-wise modelling approach with the model optimizations conducted in each of the steps.

