

Extension of the covariate clinical relevance determination in population pharmacokinetic analyses to FREM, LASSO & GA approaches



Morgane PHILIPP^{1,2}, Sylvie RETOUT^{2,3}, Simon BUATOIS³, France MENTRE¹

¹ Université Paris Cité, INSERM, IAME, UMR 1137, Paris, France

² Institut Roche, Boulogne-Billancourt, France

³ Roche Pharma Research and Early Development, Pharmaceutical Sciences, Roche Innovation Center Basel, Switzerland



CONTEXT

- Covariate clinical relevance (CCR) assessment is essential for patient dose adjustment and is a key step according to health authorities.
- Forest plots are used to illustrate parameter changes for covariate values relative to typical ones.
- CCR is inferred from the position of covariate effect ratios (CER) with their 90% confidence interval (CI90) relative to the reference area/line.
- Our previous work [1] compared the accuracy of CCR assessment of Stepwise Covariate Model (SCM), SCM+ [2] & Full Model (FM) [3]; all approaches provided satisfactory results close to those of the reference model (estimations using the simulated covariate models).
- Here, 3 alternative approaches are investigated: full random effect modelling (FREM) [4], least absolute shrinkage and selection operator (LASSO) [5] & a genetic algorithm (GA) [6]; PsN algorithms were used for FREM & LASSO; the algorithm developed by Ronchi et al. [6] was used for the GA.

OBJECTIVE

To compare the results of CCR determination by FREM, LASSO & GA with the ones by FM, SCM+ and the reference model in a proof of concept study

METHODS

Simulation study

PK model [1,7]

One compartment model (V/F)
First order absorption (ka)
Linear elimination (CL/F)

Design [1,7]

N = 383
4 phase III studies n = {1, ..., 26} &
1 phase I/II study n = {21, ..., 71}

Error model [1,7]

Combined (a and b)
→ with a the additive (fixed)
and b the proportional term

Covariate models [1,7]

Body weight (BW), age (AGE), albumin (ALB), black (BLK) race

Base model
 $CL/F_i = \mu_{CL/F} \times \left(\frac{BW_i}{70}\right)^{B_{CL/F,BW}} \times e^{\eta_{CL/F,i}}$
 $V/F_i = \mu_{V/F} \times \left(\frac{BW_i}{70}\right)^{B_{V/F,BW}} \times e^{\eta_{V/F,i}}$
 $KA_i = \mu_{KA} \times e^{\eta_{KA,i}}$
→ with μ the fixed effects and $\eta_i \sim N(0, \Omega)$ the between subject random-effects of individual i , Ω being the variance-covariance matrix, with $\beta_{par,cov}$ the effect of a covariate (cov) on a parameter (par)

Covariate model
 $CL/F_i = \mu_{CL/F} \times \left(\frac{BW_i}{70}\right)^{B_{CL/F,BW}} \times \left(\frac{AGE_i}{30}\right)^{B_{CL/F,AGE}} \times \left(\frac{ALB_i}{45}\right)^{B_{CL/F,ALB}} \times e^{\eta_{CL/F,i}}$
 $V/F_i = \mu_{V/F} \times \left(\frac{BW_i}{70}\right)^{B_{V/F,BW}} \times \left(\frac{AGE_i}{30}\right)^{B_{V/F,AGE}} \times (1 + \beta_{V/F,BLK} \times 1_{RACE_i=BLK}) \times e^{\eta_{V/F,i}}$
 $KA_i = \mu_{KA} \times e^{\eta_{KA,i}}$

Data simulation

1 dataset simulated with the **Base model** & with the **Covariate model**

Parameter estimation

Software: NONMEM 7.4

Estimation algorithm: FOCEi

SE: derived from the covariance matrix computed as $R^{-1}SR^{-1}$
(R: hessian matrix, S: cross-product gradient matrix)

Reference model fitting

Software: PsN 5.3.2

Base model & Covariate model

Covariate model building

Parsimonious covariate set (14 relations par,cov, 18 $\beta_{par,cov}$) & Saturated covariate set (21 relations par,cov, 27 $\beta_{par,cov}$)

Continuous covariates (med [min - max])	CL/F	V/F	KA	Categorical covariates category (N [%])	CL/F	V/F	KA
Body Weight (BW, kg), 69.1 [9.50-156]	B: $\beta_{CL/F,BW} = 0.939$ C: $\beta_{CL/F,BW} = 0.801$, Y, Y	B: $\beta_{CL/F,BW} = 1.066$ C: $\beta_{CL/F,BW} = 0.867$, Y, Y	N, Y	Status Non-inhibitor 195 [50%] FVIII inhibitor (INH) 194 [50%]	Y, Y	Y, Y	Y, Y
Age (AGE, years), 30.0 [1.22 - 77.00]	C: $\beta_{CL/F,AGE} = 0.127$, Y, Y	Y, Y	Y, Y	Race White, 244 [63%] Black (BLK), 31 [8%] Asian (ASN), 89 [23%] Other (OTH), 25 [6%]	Y, Y	C: $\beta_{V/F,RACE=BLK} = -0.212$, Y, Y	N, Y
Albumine (ALB, g/L), 45.0 [33.0-56.6]	C: $\beta_{ALB,BW} = -0.918$, Y, Y	Y, Y	N, Y				
Aspartate aminotransferase (AST, U/L), 23.0 [11.0-91.0]	Y, Y	N, Y	N, Y				
Bilirubin (BILI, μ mol/L), 9.0 [0.33-46.0]	Y, Y	N, Y	N, Y				

→ B, C: covariate effect simulated for the **Base** & the **Covariate model**, respectively

→ Y/N, Y/N: covariate effect tested (yes, Y) or not (No, N) on the parameter considering the **Parsimonious** or the **Saturated covariate set**

FM

Software: PsN 5.3.2
Retries = 5

SCM+

Software: PsN 5.3.2
P-cutoff = 0.05
P-forward = 0.05
P-backward = 0.01

LASSO

Software: PsN 5.3.2
Stop_t = 10

→ fit the selected covariate model with the right relation shape

GA

Software: R 4.0.5 & Perl 5.30.2
Selection pressure = 0.75
Crossover probability = 0.7
Mutation probability = 0.025

Chromosomes preserved by elitism = 10%
GA 150
150 generations
50 chromosomes
GA 50
50 generations
30 chromosomes

RESULTS

Covariates selection, CCR assessment using forest plot, OFV = $-2LL$ & BICc = $-2LL + P_p \times \ln(N) + P_e \times \ln(n_{tot})$ values, runtime
→ with LL the log likelihood, P_p the number of μ , $\beta_{par,cov}$ and $\dim(\Omega)$, P_e the number of error model parameters = 1, N the number of patients and n_{tot} the total number of observations

Forest plot generation

Software: R 4.1.2
Package: PMXForest 1.2.6
(<https://github.com/pharmetheus/PMXForest>),
PMXFrem 1.2.2
(<https://github.com/pharmetheus/PMXFrem>),
Number of samples: 1000

Possible CCR decisions illustrated on a forest plot

Clinical decision	Covariate effect ratios
Relevant (R)	→
Non-relevant significant (NRS)	→
Non-relevant non-significant (NRNS)	→
Insufficient information significant (IIS)	→
Insufficient information non-significant (IINS)	→

→ vertical reference area in light blue representing a change of $\pm 20\%$ change in parameter value compared with the typical reference individual

CONCLUSION

- FREM, LASSO & GA gave satisfactory results for CCR assessment in line with those found with FM & SCM+.
- FREM & FM gave very comparable results, making both methods very suitable for CCR assessment.
- GA150 selected more parsimonious models than GA50 & LASSO and resulted in better BIC; however the number of models generated is substantial and requires a high run time; if computing resources are not an issue, GA150 may be preferable.
- Between full modelling methods and covariate selection methods, the former have the benefits to get a CCR evaluation of all relations while the latter provide a parsimonious model suitable for prediction; the best choice depend on one's objective.
- Further evaluations are necessary to enhance the robustness of those results by using simulation frameworks with increased complexity in model structures, expanded covariate sets, and a larger number of simulated datasets.

References

[1] Philipp et al. J. Pharmacokinet. Pharmacodyn. 2024; [2] Svensson et al. CPT Pharmacomet. Syst. Pharmacol. 2022; [3] Xu et al. Br. J. Clin. Pharmacol. 2018; [4] Yngman et al. CPT Pharmacomet. Syst. Pharmacol. 2022; [5] Ribbing et al. J. Pharmacokinet. Pharmacodyn. 2007; [6] Ronchi et al. Pharmacokinet. Pharmacodyn. 2023; [7] Retout et al. Clin. Pharmacokinet. 2020;

RESULTS

Covariates selection & CCR assessment

Simulation case: Base model							Simulation case: Covariate model						
Ref model	FM	FREM	SCM+	LASSO	GA 150	GA 50	Ref model	FM	FREM	SCM+	LASSO	GA 150	GA 50
Parsimonious covariate set													
CL/F, BW			/						/				
CL/F, AGE			/						/				
CL/F, ALB			/						/				
CL/F, AST			/						/				
CL/F, BILI			/						/				
CL/F, RACE			/						/				
CL/F, INH			/						/				
V/F, BW			/						/				
V/F, AGE			/						/				
V/F, ALB			/						/				
V/F, RACE			/						/				
V/F, INH			/						/				
KA, AGE			/						/				
KA, INH			/						/				
Saturated covariate set													
CL/F, BW													
CL/F, AGE													
CL/F, ALB													
CL/F, AST													
CL/F, BILI													
CL/F, RACE													
CL/F, INH													
V/F, BW													
V/F, AGE													
V/F, ALB													
V/F, AST													
V/F, BILI													
V/F, RACE													
V/F, INH													
KA, BW													
KA, AGE													
KA, ALB													
KA, AST													
KA, BILI													
KA, RACE													
KA, INH													

R

NRS

NRNS

IIS

IINS

NSEL

NIM

/

NP


NSEL: not selected, NIM: not in the model, NP: not performed

Covariate selection

- Covariate selection approaches (i.e. SCM+, LASSO & GA) selected all simulated relations
- Few additional relationships were retained, with a larger number using the saturated covariate set and with SCM+ being the more restrictive in selection

CCR assessment

- $\beta_{par,cov} = 0$: all covariates modelling methods led to consistent CCR assessment and in line with those found with the reference model
→ BW on CL/F & V/F: **R**; AGE on CL/F & V/F & RACE on V/F: **IIS**; ALB on V/F: **NRS**
- $\beta_{par,cov} = 0$: none of covariates modelling methods conclude to the R of one of these relations
 - Full modelling approaches (i.e. FM & FREM): NR or with II to conclude
 - Covariate selection approaches: mainly NSEL and the few spurious relations retained NR or with II to conclude



OFV, BICc & runtime

	Simulation case: Base model						Simulation case: Covariate model							
	Ref model	FM	FREM*	SCM+	LASSO	GA 150	GA 50	Ref model	FM	FREM*	SCM+	LASSO	GA 150	GA 50
Parsimonious covariate set														
OFV	24 135	24 110	/	24 126	24 121	24 126	24 126	24 056	24 038	/	24 056	24 039	24 044	24 050
P _r	8	24	/	9	16	9	11	14	24	/	14	21	16	19
BICc	24 191	24 261	/	24 188	24 225	24 188	24 200	24 148	24 189	/	24 148	24 172	24 148	24 172
Runtime (h)	0.04	0.44	/	0.66	3.31	5.99	2.15	0.04	0.44	/	0.81	5.38	9.89	2.32
Saturated covariate set														
OFV	24 135	24 103	19 786*	24 126	24 119	24 121	24 114	24 056	24 032	19 715*	24 048	24 046	24 044	24 042
P _r	8	33	90	9	17	11	15	14	33	20 259*	15	20	16	22
BICc	24 191	24 308	20 330*	24 188	24 229	24 195	24 212	24 148	24 237	20 259*	24 146	24 173	24 148	24 181
Runtime (h)	0.04	0.44	2.05	1.55	4.51	5.99	2.09	0.04	0.44	2.18	2.09	6.33	10.73	3.71

*FREM OFV & BICc are not comparable with other methods due to differences regarding the dataset (covariates treated as observations), the way of coding covariate effects (as random-effects) and the Ω matrix (omega block for FREM vs diagonal matrix for the other methods)

OFV & BICc

- FM → best OFV in all cases, but a worse BIC due to the very large number of $\beta_{par,cov}$ to be estimated
- FREM → OFV and BICc not comparable with other methods, see *
- SCM+ → best BICc in all cases, with values lower than or equal to the ones of ref model
- GA 50 → results very close to the ones of SCM+ in terms of BICc (from 0 to 7 points difference)
- LASSO & GA 150 → slightly higher BICc (9 to 38 points compared to ref model) due to less parsimonious models selected

Runtime

- FM showed the best runtime; only SCM+ can compete with the runtime of the full modelling methods
- GA 50 & 150: high runtime due to the large number of models to run (max of 1500/7500 for GA50/150)
- LASSO: runtime depends mainly on the cross-validation number of iterations (10 models ran at each iteration)