

Debra van Asten^{1,2}, Frédéric Gaspar^{1,3,4}, Anne Ravix¹, Jean Terrier^{5,6,7}, Pauline Gosselin^{5,6}, Pierre Fontana^{6,8}, Youssef Daali^{6,7}, Jean- Luc Reny^{5,6},
Chantal Csajka^{1,3,4*}, Monia Guidi^{1,4,9*}

Introduction

- ## Methods & Data

- Ticagrelor and AR-C124910XX data were modelled simultaneously, via a classic stepwise population PK (popPK) approach (Monolix 2021), assuming **linear and irreversible biotransformation** between the parent drug to the metabolite.
- Rich **dried bloodspot (DBS)** data was collected from 103 hospitalized patients* who received a dose of 90 mg ticagrelor and reached steady state prior to the start of the **OptimAT study**¹.
- Tested covariates included demographic (e.g. body weight, age and gender) and genetic factors (CYP3A4/5 and P-glycoprotein (P-gp) phenotypic activity). Individual phenotypes were obtained via the **Geneva Cocktail approach** for determination of CYP and P-gp activities².

Results

```

graph LR
    Dose[Dose] -- "T_Lag  
D_1" --> Tigacrelor_Vcentral((Tigacrelor  
Vcentral))
    Tigacrelor_Vperipheral((Tigacrelor  
Vperipheral)) -- "K_21" --> Tigacrelor_Vcentral
    Tigacrelor_Vcentral -- "K_12" --> Tigacrelor_Vperipheral
    Tigacrelor_Vcentral -- "CL_ptot * (F_tr)" --> AR_C124910XX_Vcentral((AR-C124910XX  
Vcentral))
    AR_C124910XX_Vcentral -- "K_43" --> AR_C124910XX_Vperipheral((AR-C124910XX  
Vperipheral))
    AR_C124910XX_Vperipheral -- "K_34" --> AR_C124910XX_Vcentral
    Tigacrelor_Vcentral -- "CL_ptot * F_CL" --> Elim1[Elimination]
    AR_C124910XX_Vcentral -- "CL_m" --> Elim2[Elimination]
  
```

Label	Estimate (RSE%)	Estimate (RSE%)
	Population parameters (θ)	Inter-individual variability (CV(%))
Parent		
D ₁ (h)	1.07 (7.3)	65 (11.1)
T _{Lag} (h)	0.34 (7.4)	66 (9.3)
CL _{ptot} (L/h)	18.9 (4.4)	45 (7.4)
V _{pc} (L)	118 (6.5)	57 (9.7)
V _{pp} (L)	326 (11.0)	125 (9.0)
Q _p (L/h)	59.8 (5.1)	-
F _{CLp}	0.36 (1.5)	-
Metabolite		
CL _m (L/h)	21.6 (3.7)	36 (7.7)
Age on CL _m	-1.11 (18.4)	-
V _{mc} (L)	1	-
V _{mp} (L)	41.6 (13.9)	119 (13)
Q _m (L/h)	29.2 (2.7)	-
Correlations		
V _{pc} and CL _p	0.73	11.4
Residual variability (CV(%))		
Parent: Proportional	13.1 (3.3)	
Metabolite: proportional	10.0 (3.1)	

Main findings

- A 2-compartmental model structure with **zero-order absorption and a lag-time** accurately characterized OptimAT data.
- An inverse covariate relationship between **CL_m and age** was included in the model. An increase of 10-years in age produces a 15% relative decrease (3.2 L/h) in CL of the metabolite.
- No significant associations could be identified between PK model parameters and other available covariates, **including CYP3A4/5 and P-gp phenotypic activity**.

Conclusion

- This study confirmed the substantial variability in the PK profile of ticagrelor and its metabolite in DBS samples.
- Further research is needed to confirm the influence and study the clinical significance of the large IIV and the relationship between CL of the metabolite and age.

1. Antithrombotics' Therapeutic Optimization in Hospitalized Patients Using Physiologically- and Population-based Pharmacokinetic Modeling.
2. Bosilkovska, M., et al., Geneva cocktail for cytochrome p450 and P-glycoprotein activity assessment using dried blood spots. *Clin Pharmacol Ther.* 2014. 96(3): p. 349-59.