


Replacing transit compartments with the analytical solution of a partial differential equation in malaria liver-stage modeling

Luca Barberi, Sam Jones & Nathalie Gobeau
MMV Medicines for Malaria Venture

1. Background

After infection by a mosquito bite, malaria *P. falciparum* parasites go into the liver where they stay and replicate for over one week before invading the blood.



Figures from Wikimedia Commons.

2. Model of the liver stage

Parasite ageing in the liver can be modeled using a Partial Differential Equation (PDE) [1]. This is mathematically equivalent to a fluid (parasites) flowing through a pipe (age). In addition to ageing, parasites also replicate and are killed by drugs.

$$\frac{\partial p_L}{\partial t} + \frac{\partial p_L}{\partial a} = p_L(GR_L - Kill_L) \quad P_L(t) = \int_0^{T_L} p_L(a, t) da$$

Definitions in the **List of symbols** below.


3. Solving the model: transit compartments

The PDE can be numerically approximated by a set of ODEs, mathematically equivalent to transit compartments.

$$\begin{aligned} \frac{d}{dt} p_L^1(t) &= F_{inf} P_{spz} INPUT - k_{tr} p_L^1 + (GR_L - Kill_L) p_L^1 \\ \frac{d}{dt} p_L^i(t) &= k_{tr} (p_L^{i-1} - p_L^i) + (GR_L - Kill_L) p_L^i \\ \frac{d}{dt} p_L^{N_{cpt}}(t) &= k_{tr} p_L^{N_{cpt}-1} - (GR_L - Kill_L) p_L^{N_{cpt}} \end{aligned}$$

Where $k_{tr} = N_{cpt}/T_L$ and $i = 2 \dots N_{cpt} - 1$. Parasites from different bites are passed as “input doses” to p_L^1 .

Example: 1 bite at $t = 0$, $GR_L = Kill_L = 0$, $T_L = 6$ days, $F_{inf} P_{spz} = 1$. Slope of liver escape determined by N_{cpt} (numeric diffusion).



4. Solving the model: analytical solution


The PDE can be solved analytically.

$$p_L(a, t) = F_{inf} P_{spz} \sum_{n=1}^{N_{bites}} p_L(a - (t - t_n), t = t_n) e^{J_n}$$

$$\frac{dJ_n}{dt} = \begin{cases} 0, & t < t_n \\ GR_L - Kill_L, & t \geq t_n \end{cases}$$

We assume that the initial distribution of parasite ages $p_L(a, t = t_n)$ is the same for all bites. To perform simulations, we assume this is a (narrow) Gaussian, $p_L(a, t = t_n) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{1}{2}(\frac{a}{\sigma})^2}$.

Example: 1 bite at $t = 0$, $GR_L = Kill_L = 0$, $T_L = 6$ days, $F_{inf} P_{spz} = 1$. Slope of liver escape determined by σ .



5. Conclusions

- The analytical solution does not require to solve a possibly high number (N_{cpt}) of ODEs.
- The slope of liver escape is simply determined by σ , not numeric diffusion (via N_{cpt}).
- With limited computational effort, the analytical solution allows to track the dynamics of parasites from individual bites, useful to introduce “cure thresholds” (i.e. number of parasites from the same bite can be either ≥ 1 or 0).

List of symbols

p_L = parasite density in liver
 a = parasite age
 T_L = avg. time spent in liver
 GR_L = growth rate in liver
 $Kill_L$ = drug kill rate in liver
 P_L = n. parasites in liver
 $F_{inf} P_{spz}$ = n. parasites transferred to liver from single bite
 t_n = time of n^{th} bite

References

[1] C. Barcelo, S. Jones, P. J. Lowe, J. J. Möhrle, N. Gobeau, and M. H. Cherkaoui-Rbati, *In Preparation*.

