

Efficient and Informed Model Development for NLME Models with Poor Parameter Identifiability

Karsten Kuritz, Daniel Kaschek, Daniel Lill, IntiQuan AG, Basel, Switzerland
Nathalie Gobeau, MMV Medicines for Malaria Venture, Geneva, Switzerland

IntiQuan⁺
A ProductLifeGroup Company

Problem

- Poor parameter identifiability can lead to convergence issues and unreliable prediction due to inflated RSEs.
- This slows model development and undermines confidence in simulation outcomes.


Approach

- Use of dMod¹ in IQRtools as deterministic framework combining trust-region optimization with profile likelihood analysis to detect and resolve non-identifiabilities.
- Implemented using analytically derived gradients and Hessians for high precision.
- Inter-individual variability is modeled using a penalized likelihood, enabling efficient and robust estimation.

Impact

- In a malaria PK/PD model, our approach identified a poorly identifiable Hill coefficient.
- Profile likelihood guided parameter fixing, resulting in a stable, interpretable model.
- The framework improves model reliability, supports informed decisions, and reduces development time.

Hill = 23.7 (947% RSE)



Application Example: Malaria PD Modeling


In malaria PK/PD modeling, parasite killing is described by a .model linking drug concentration to the parasite elimination rate.

$$\frac{dP}{dt} = \left(GR - Emax \frac{C c^{hill}}{C c^{hill} + EC_{50}^{hill}} \right) \cdot P$$



Example 1: Preclinical study

Objective:

Characterize the PK/PD relationship in SCID mice infected with *P. falciparum* to support human dose prediction.



- Hill coefficient **poorly identified** (hill = 23.7, 947% RSE) due to limited data support.
- In trial simulations this causes instability **from extreme parameter values**.
- Profile likelihood analysis reveals a **right-open shape** for hill.




Parameter	Hill estimated	Hill fixed
GR	0.03 (FIX)	0.03 (FIX)
EMAX	0.24 (FIX)	0.24 (FIX)
EC50	0.00699 (13%)	0.007 (12%)
hill	23.7 (950%)	20 (FIX)
omega(EMAX)	0.1 (FIX)	0.1 (FIX)
omega(EC50)	0.3 (FIX)	0.3 (FIX)
error_ADD1	0.75 (7.4%)	0.75 (7.4%)
OBJ	245	245
BIC	265	260

Fixing hill to 20 improved model identifiability while accounting for variability in EC50.



Example 2: Clinical Phase 1b Study⁴

Objective:

Characterize the PK/PD relationship between pyronaridine blood concentrations and *P. falciparum* parasitemia in healthy volunteers with induced blood-stage malaria, to inform dose prediction in African children.



Multi-start optimization revealed two optima with distinct objective function values. Profile likelihoods show well-behaved, **symmetric shapes** for all parameters.



Parameter estimates are **consistent** across tools (vs. Monolix), supporting **model robustness**.

Parameter	dMod	Monolix
GR	0.0871 (4.1%)	0.0856 (3.9%)
EMAX	0.204 (7.3%)	0.201 (3.9%)
EC50	0.0058 (8.3%)	0.00575 (11%)
hill	6.3 (11%)	6.62 (2.4%)
omega(GR)	0.1 (FIX)	0.1 (FIX)
omega(EMAX)	0.2 (FIX)	0.0996 (29%)
omega(EC50)	0.2 (FIX)	0.273 (29%)
error_ADD1	1.04 (3.2%)	1.02 (3.4%)

IQRtools integrates gradient-based optimization and profile likelihoods with population modeling

IQRmodel syntax for consistent model description

- Versatile modeling syntax** compatible with formats of typically used tools in QSP and PMX.
- A **PK model library** is available in the documentation material.

```
***** MODEL NAME
Linear 1-compartmental PK model

***** MODEL NOTES
Linear 1-compartmental PK model with initial single dose.


***** MODEL STATES
d/dt(Ac) = -CL/Vc*Ac + INPUT1
Ac(0) = 0

***** MODEL PARAMETERS
CL = 20 # (L/hour) Clearance
Vc = 32 # (L) Central volume

***** MODEL VARIABLES
Cc = Ac/Vc * 1000 # (ng/mL) Plasma concentration
```


Highly efficient parameter estimation

- State-of-the-art optimization using analytically derived **sensitivity equations**,
- Deterministic optimization** combined with,
- Global sampling** of initial guesses.




Identification of structural and practical non-identifiability

- Nonlinear likelihood** function shape is not captured by the **Fisher information matrix**.
- Likelihood profiling** characterizes the parameter identifiability and characterization of the **parameter identifiability**.



$$PL_j(p) = \max_{\theta \in \{\theta | \theta_j = p\}} \ln L(Y|\theta)$$



How we deal with variability of parameters between virtual patients

- Maximum a posteriori estimation / penalized **maximum likelihood** estimation.

$$f(\mu)f(x|\mu) = \pi(\mu)L(\mu) = \frac{1}{\sqrt{2\pi}\sigma_m} \exp\left(-\frac{1}{2}\left(\frac{\mu - \mu_0}{\sigma_m}\right)^2\right) \prod_{j=1}^n \frac{1}{\sqrt{2\pi}\sigma_v} \exp\left(-\frac{1}{2}\left(\frac{x_j - \mu}{\sigma_v}\right)^2\right)$$

References:

- Kaschek D, Mader W, Fehling-Kaschek M, Rosenblatt M, Timmer J. Dynamic modeling, parameter estimation, and uncertainty analysis in R. J Stat Softw. 2019;88(10):1–32. doi:10.18637/jss.v088.i10
- Raue A, Schilling M, Bachmann J, et al. Lessons learned from quantitative dynamical modeling in systems biology. PLoS One. 2013;8(9):e74335. doi:10.1371/journal.pone.0074335
- Raue A, Kreutz C, Maiwald T, Klingmüller U, Timmer J. Addressing parameter identifiability by model-based experimentation. IET Syst Biol. 2011;5(2):120–130. doi:10.1049/iet-syb.2010.0030
- Barber BE, Webster R, Potter AJ, et al. Characterising the blood-stage antimalarial activity of pyronaridine in healthy volunteers experimentally infected with Plasmodium falciparum. Int J Antimicrob Agents. 2024;64:107196. doi:10.1016/j.ijantimicag.2023.107196

MMV
Medicines for Malaria Venture

IntiQuan⁺
A ProductLifeGroup Company

