



Population pharmacokinetics of high-dose isoniazid in children receiving 3-month weekly isoniazid and rifapentine (3HP) for TB preventive treatment

Ruvarashe J Madzime^{1*}, Louvina E van der Laan², Lufina Tsirizani¹, Lubbe Wiesner¹, Belen P Solans³, Rada Savic³, Kelly Dooley⁴, Shaun Barnabas⁴, Ingrid Courtney², Avy Violari⁶, Paolo Denti¹, Anneke C Hesselning²

¹ Division of Clinical Pharmacology, Department of Medicine, University of Cape Town, South Africa; ² Desmond Tutu TB Centre, Department of Paediatrics and Child Health, Faculty of Medicine and Health Sciences, Stellenbosch University, South Africa, ³ University of San Francisco School of Pharmacy, California, USA, ⁴ Division of Infectious Diseases, Department of Medicine, Vanderbilt University Medical Center, ⁵ FAMCRU, Department of Paediatrics and Child Health Faculty of Medicine and Health Sciences, Stellenbosch University, South Africa, ⁶ Baragwaneth, Perinatal HIV Research Unit (PHRU), Wits Health Consortium, Johannesburg, South Africa



*Email: mdzruv004@myuct.ac.za

Background and objectives

Children are at an increased risk of developing severe forms of TB disease upon TB infection, if left untreated¹.

WHO recommends **3HP** (once weekly isoniazid and rifapentine for 3 months) regimen for the treatment of TB infection in adults (more recently in children)².

Isoniazid pharmacokinetics is not well characterized in children under the age of 2 years.

Children under 2 years of age are still **maturing** and developmental pharmacology impacts the pharmacokinetics of a drug.

Isoniazid clearance is impacted by the polymorphic **NAT2 enzyme**, and this effect is not well characterized in infants below 2 years of age.

Previous studies have reported **lower bioavailability** of isoniazid, especially in younger children³.

We determined the pharmacokinetics of **high-dose isoniazid (15-25mg/kg)** given once weekly with rifapentine for 3 months for prevention of TB disease in children.

Methods

This was part of the **Tuberculosis Trials Consortium Study 35**, and it involved **children in South Africa aged 0 – 12 years**, with or without HIV and with exposure to drug-susceptible TB or *M.tb* infection but with no active TB disease.




Figure 1. Study protocol

Isoniazid plasma concentrations were assayed using **LC-MS** with a **LLOQ of 0.105mg/L**.

We started analysis from a literature model that was developed for **South African children on first line TB treatment (10 mg/kg)**³. The model was updated in **NONMEM** version 7.5 using **FOCE-I**.

Below Limit of Quantification (**BLQ**) values were handled using **M7+** method⁴. Allometric scaling was tested on disposition parameters.

A **mixture model**⁵ was implemented to **impute the NAT2 acetylator** status in the population.

The area under the curve (AUC) was calculated using the final model.

Results

Table 1. Participants characteristics

Variable	Age cohorts				
	Cohort 1 ≥ 4 to ≤ 12y N = 9	Cohort 2 ≥ 2 to < 4y N = 6	Cohort 3 ≥ 1 to < 2y N = 19	Cohort 4 0 to < 1y N = 19	Overall N = 53
Age [years]	11.2 (7.3, 12.9)	2.8 (2.6, 3.1)	1.5 (1.1, 2.0)	5.76 (0.84, 11.28) months	1.3 (0.1, 12.9)
Weight [kg]	27 (18, 39)	15 (12, 15)	9 (8, 17)	7 (4, 10)	9 (4, 39)
Height [cm]	127 (112, 152)	91 (88, 96)	78 (73, 85)	66 (53, 75)	77 (53, 152)
Living with HIV	7 (78%)	0 (0%)	0 (0%)	0 (0%)	7 (13%)
Sex					
Female	4 (44%)	2 (33%)	7 (37%)	9 (47%)	22 (42%)
Male	5 (56%)	4 (67%)	12 (63%)	10 (53%)	31 (58%)
Race					
Black	2 (22%)	4 (67%)	12 (63%)	11 (58%)	29 (55%)
Mixed	7 (78%)	2 (33%)	7 (37%)	8 (42%)	24 (45%)
Median (min, max) or Frequency (%)					




Figure 2. Final pharmacokinetic model. ktr: transit time; NN: number of transit compartments; ka: absorption rate constant; Q: intercompartmental clearance; CL: clearance; V: central compartment volume of distribution; V3: peripheral volume of distribution. WT represents allometric scaling with total body weight on CL(0.75), Q(0.75), V(1), and V3(1). AGE represents the maturation on clearance. The **mixture model** describes the differences of NAT2 acetylation within the population.

Results

Isoniazid pharmacokinetics were best described by a 2 compartment disposition model (Figure 2).

Of the total available observations, approximately **36% were BLQ** (mostly at pre-dose).

Disposition parameters (CL, Q, V and V3) were **allometrically scaled** using **total body weight**.

Mixture model was fixed to **2 subpopulations** (fast and intermediate and slow populations) for model stability (Figure 2).

Maturation function indicated that **50% maturation** is attained at approximately **1 month** and **≥98% maturation** at approximately **18 months**.

There were **no grade 3 or higher** adverse events – **good safety outcomes**.




Figure 3. Visual predictive checks (VPC) of the model showing the 95% confidence interval of the model predicted and observed data. The y-axis shows the observed isoniazid concentrations in mg/L and the x-axis shows the time after last dose in hours. **A** – VPC is stratified by model imputed acetylator status. **B** – VPC is stratified by the age cohorts.

4




Figure 4. Jitter plot for isoniazid AUCs, y-axis shows isoniazid AUC₀₋₂₄ in mg·h/L and the x-axis shows the 4 age cohorts in the study. The bars represent the median and the 25th and 75th percentiles. The **dark purple bars** represent the **fast and intermediate acetylator** group, and the **light purple bars** represent the **slow acetylator** group. The green dashed line shows this study's (**study 35**) median of **31 mg·h/L**. The red dashed line shows the **Shine trial**⁶ median of **9.7 mg·h/L**. The blue dashed line shows the median AUC of **78 mg·h/L** from a 3HP trial⁷ conducted in Thailand.

5




Figure 5. Maturation of clearance. The y-axis shows the maturation in percentage. The x-axis is the post-natal age in months, and the vertical lines at the bottom represent the ages in our dataset.

Conclusions

The **model (high-dose isoniazid 15-25 mg/kg)** was in line with **literature findings (first line TB treatment – 10mg/kg)**.

Identified an **age effect on clearance** – maturation function in line with literature values.

Identified **NAT2 bimodal effect** on clearance.

Contrary to previous findings, age did not contribute to a change in bioavailability, which may be due to the use of the same formulation in our study.

Observed a **linear dose-exposure** relationship (Figure 4), which is in line with literature findings.

High exposures did not lead to adverse events, safety outcomes were good.

Findings from this model informed recent WHO guidelines.

Acknowledgements



Presentation of this poster and participation in the 33rd PAGE meeting was made possible thanks to a **PAGE scholarship**. Financial support was based on predefined criteria, and the merit of each application was assessed by a selection committee.

References

- Marais BJ, Gie RP, Schaaf HS, et al. The natural history of childhood intra-thoracic tuberculosis: a critical review of literature from the pre-chemotherapy era. Int J Tuberc Lung Dis. 2004;8(4):392-402.
- World Health Organization. WHO Consolidated Guidelines on Tuberculosis. Module 1: Prevention - Tuberculosis Preventive Treatment. Second.; 2024. Accessed February 4, 2025. <https://www.who.int/publications/i/item/9789240096196>
- Galileya LT, Wasmann RE, Chabala C, et al. Evaluating pediatric tuberculosis dosing guidelines: a model-based individual data pooled analysis. PLoS Med. 2023;20(11):e1004303. doi:10.1371/journal.pmed.1004303
- Wijk M, Wasmann R, Jacobson K.R, Svensson E. M, Denti P. A Pragmatic Approach to Handling Censored Data Below the Lower Limit of Quantification in Pharmacokinetic Modeling. CPT: Pharmacometrics & Systems Pharmacology; 2025.
- Ette EI, Williams Paul J. Pharmacometrics: The Science of Quantitative Pharmacology. First. (Ette EI, Williams Paul J, eds.). John Wiley & Sons, Inc; 2007.
- Chabala C, Turkova A, Thomason MJ, et al. Shorter treatment for minimal tuberculosis (TB) in children (SHINE): a study protocol for a randomised controlled trial. Trials. 2018;19(1):237. doi:10.1186/s13063-018-2608-5
- Phaisal W, Jantarabenjakul W, Wacharachaisurapol N, et al. Pharmacokinetics of isoniazid and rifapentine in young pediatric patients with latent tuberculosis infection. Int J Infect Dis. 2022;122:725-732. doi:10.1016/j.ijid.2022.07.040