


POPULATION PHARMACOKINETIC MODELLING TO CHARACTERIZE THE EFFECT OF CHRONIC KIDNEY DISEASE ON TENOFOVIR EXPOSURE AFTER TENOFOVIR ALAFENAMIDE ADMINISTRATION

Paul THOUEILLE¹, Susana ALVES SALDANHA¹, Vincent DESFONTAINE¹, Katharina KUSEJKO^{2,3}, Perrine COURLET¹, Pascal ANDRE¹, Matthias CAVASSINI⁴, Laurent A. DECOSTERD¹, Thierry BUCLIN¹, Monia GUIDI^{1,5,6},
The Swiss HIV Cohort Study

Contact: Paul.Thouelle@chuv.ch

Introduction


- Tenofovir is an **NRTI** active in preventing and treating HIV-1 infection, and treating HBV infection.
- Tenofovir plasma levels are associated with **kidney tubular damage** and, in the long term, **bone demineralization**.
- Two prodrugs available** (*dosing 1x/day*):



Objectives

- To characterize the **concentration-time profile of tenofovir and its associated variability** in people living with HIV (PLWH) under TAF in a *real-life* setting.
- To assess the **impact of chronic kidney disease (CKD)** on tenofovir exposure.

Methods



Results


Parameters	Final model				Bootstrap (n = 2000 samples)			
	Estimate	RSE (%)	BSV (%)	RSE (%)	Median	CI _{95%}	BSV (%)	CI _{95%}
k_a (h ⁻¹)	2 FIX	-			2 FIX	-		
V_{TFV} (L)	2390	9			2397	2037 - 2891		
CL_{TFV} (L/h)	42.2	2			42.3	40.1 - 44.3		
θ_{CLCR}	0.707	10			0.703	0.549 - 0.837		
θ_{Age}	-0.244	30			-0.242	-0.398 to -0.099		
$\theta_{Black \text{ or } Hispano \text{ American}}$	0.119	23			0.120	0.068 - 0.175		
$\theta_{Potent \text{ P-gp inhibitors}}$	-0.121	39			-0.119	-0.202 to 0		
F (%)	1 FIX	-	21.1	6	1 FIX	-	20.9	18.4 - 23.5
$\theta_{Cobicistat}$	1.15	6			1.16	1.02 - 1.28		
σ_{add} (nmol/mL)	0.0099	5			0.0099	0.0088 - 0.0110		

RSE: Relative standard error BSV: Between-subject variability CI_{95%} 95% confidence interval

Final model: θ_X , effect of the X covariate on the typical value (TV) of CL_{TFV}

TAF 10 mg: $F = (1 + \theta_{cobicistat})$

$$TVCL_{TFV} = CL_{TFV} \times \left(1 + \frac{\theta_{CLCR} \times (CL_{CR} - CL_{CR-ref})}{CL_{CR-ref}}\right) \times \left(1 + \frac{\theta_{Age} \times (Age - Age_{median})}{Age_{median}}\right) \times (1 + \theta_{Black \text{ or } Hispano \text{ American}}) \times (1 + \theta_{Potent \text{ P-gp inhibitors}})$$




Simulated PK profiles for tenofovir after 25 mg TAF administration in individuals with different CKD stages, with the 2.5%, 25%, 50% (median), 75% and 97.5% percentiles (red solid lines), $PI_{50\%}$ and $PI_{95\%}$ (shaded areas).

Covariates effect


- A full model incorporating all covariates explained 59% of the BSV on F.
- A reduced model with CL_{CR} alone as covariate explained 53% of the BSV on F.
- Only the effect of CL_{CR} on CL_{TFV} was considered clinically relevant for simulations of PK profiles as a function of different CKD stages.**

Schematic representation of the PK models

Model with TAF and tenofovir



Model with tenofovir only



k_{12} : absorption rate constant from depot to the TAF compartment; k_a : first-order absorption rate constant of tenofovir; k_{23} : rate constant from TAF compartment to the tenofovir compartment; CL_{TAF} : apparent clearance of TAF; V_{TAF} : apparent volume of distribution of TAF; CL_{TFV} : apparent clearance of tenofovir; V_{TFV} : apparent volume of distribution of tenofovir; k_e : tenofovir elimination rate constant.

Conclusions

- PLWH with stage 3 and stage 4 CKD under TAF achieve similar tenofovir exposure to those with normal kidney function receiving TDF.
- Evidence²⁻⁶ suggests that TAF is more permissive than TDF with respect to missed doses or longer dosing intervals, and that **PLWH with stage 3 or 4 CKD could potentially receive TAF once every 2 days or 3 days**, respectively, in order to maintain adequate circulating levels of tenofovir while preserving sufficient tenofovir-diphosphate exposure.

References

- Courlet P, et al. Clinical Mass Spectrometry. 2018
- Thurman AR, et al. eClinicalMedicine. 2021
- Podany AT, et al. AIDS. 2018
- Yager JL, et al. AIDS. 2021
- Brooks KM, et al. The Journal of Antimicrobial Chemotherapy. 2020
- Dumond JB, et al. Antiviral Therapy. 2019