

A Bayesian Optimization Procedure for the Automated Determination of Optimal Limited Sampling Strategies

A. Janssen¹, F.C. Bennis², M.H. Cnossen³, and R.A.A. Mathôt¹ for the OPTI-CLOT study group and SYMPHONY consortium

¹ Department of Clinical Pharmacology, Hospital Pharmacy, Amsterdam University Medical Center, Amsterdam, The Netherlands.

² Quantitative Data Analytics Group, Department of Computer Science, VU Amsterdam, Amsterdam, The Netherlands

³ Department of Paediatric Haematology, Erasmus MC Sophia Children's Hospital, Erasmus University Medical Center Rotterdam, The Netherlands

INTRODUCTION

Limited sampling strategies

- PK-guided dosing is a common approach for individualizing prophylactic treatment of haemophilia A patients.
- Patient blood samples are collected to individualize PK parameter estimates from prior population-PK models.
- Limited sampling strategies (LSSs) are often obtained via simulation experiments^{1,2,3} where several hypotheses concerning the number and timing of samples are evaluated.


Opportunities for improvement

- Generally only convenient sampling times (e.g. in 24h intervals) are considered.
- No information on contribution of each unique sample on accuracy.
- Differences in sub-populations can only be analysed *a priori*.
- Analysis involves a lot of manual work.
- Often only the accuracy of predictions is evaluated, while its uncertainty might be more informative in practice.

Bayesian Optimization

- Bayesian optimization (BO) is a method for the efficient optimization of computationally expensive black-box functions.
- Generally a Gaussian Process ($\mathcal{GP}$) is used. A $\mathcal{GP}$ represents a distribution over functions.
- Observations are added in order to update the posterior distribution over these functions.
- An acquisition function is then used to select the most informative point to sample next.
- The resulting function $f(t)$ represents the improvement in the objective over time t .

METHOD




USE-CASE: HAEMOPHILIA A

- Compare BO-LSS to prior LSSs from the literature.
- We use posterior variance of $p(\eta | y)$ as proxy for prediction uncertainty.

APPROACH


- Simulate patients from PK model of literature study and run BO.
- For each sample, fit $\mathcal{GP}$ to $p(\eta | y)$ over t , take $\max(\sum_i \sigma^2[\eta_i])$, and repeat.

STANDARD HALF-LIFE FVIII



- BO-LSS: samples at $t = 1$ and two at 16h post-infusion.
- Literature: samples at $t = 4, 24$, and 48h post-infusion¹.

EXTENDED HALF-LIFE FVIII



- BO-LSS: two samples at $t = 1$ and one at 48h post-infusion.
- Literature: samples at $t = 1$ and two at 24, 48, or 72h post-infusion².

Table 1. Accuracy of BO-LSS versus literature LSS

Approach	Overall accuracy (RMSE)		Residual uncertainty (σ)				Peak interval accuracy (RMSE)		Trough interval accuracy (RMSE)	
	SHL	EHL	SHL		EHL		SHL (0 - 4h)	EHL (0 - 4h)	SHL (36 - 48h)	EHL (72 - 120h)
			CL	Vd	CL	Vd				
Literature	0.048	0.042	0.17	0.14	0.086	0.11	0.085	0.082	0.027	0.014
BO	0.037	0.042	0.15	0.084	0.10	0.086	0.056	0.077	0.021	0.014

BO: Bayesian optimization, RMSE: root mean squared error, SHL: standard half-life, EHL: extended half-life, CL: clearance, Vd: volume of distribution, σ : standard deviation

REFERENCES

- [1] Bjorkman S. *Haemophilia*. 2010;16(4):597-605.
- [2] McEneny-King, Alanna, et al. *Haemophilia* 27.3 (2021): 408-416.
- [3] Preijers T., et al. *Haemophilia*, 22 (2016), pp. 100-101