

# Influence of immunization on the pharmacokinetics of monoclonal antibodies : anti-drug antibody-mediated drug disposition (ADAMDD) of rituximab (RTX) in mice

David TERNANT<sup>1</sup>, Hervé WATIER<sup>2</sup>, Céline DESVIGNES<sup>1</sup>, Nicolas AZZOPARDI<sup>3</sup>, Alain LEPAPE<sup>4</sup>, Valérie GOUILLEUX\_GRUART<sup>2</sup>, Gilles PAINTAUD<sup>1</sup>

<sup>1</sup>Université de Tours, EA 4245 «Transplantation, Immunology, Inflammation», Tours, France; CHRU de Tours, service de Pharmacologie Médicale, Tours, France. <sup>2</sup>EA 7501 « Groupe Innovation et Ciblage Cellulaire », Tours, France; CHRU de Tours, service d’Immunologie, Tours, France. <sup>3</sup>EA 7501 « Groupe Innovation et Ciblage Cellulaire », Tours, France. <sup>4</sup>CNRS UPS44, CIPA, PHENOMIN-TAAM, Orléans, France.

## INTRODUCTION

- Immunisation against monoclonal antibodies (mAbs) lead to the development of **anti-drug antibodies (ADA)** responsible of **increase in mAb clearance** by approximately 1.5 fold [1].
- Big challenge in human mAb therapy
- Joint kinetics of both mAbs and anti-mAb ADA in previous works suggested:
  - nonlinear elimination of mAbs [2]
  - mechanism of mAb elimination similar to target-mediated drug disposition (TMDD) [3]
- Objectives.** To describe the kinetics of immunisation and interactions of rituximab (RTX), an anti-CD20 mAb, and anti-drug antibodies (ADA) in mice using TMDD-derived models [4,5] defined as ADA-mediated drug disposition (ADAMDD)

## METHODS

- 129 male mice** of 9 different lines administered a single 50 µg dose of rituximab in tail vein
- 829 blood samples were collected at hours 1, 6 and 24, and days 3, 7, 15 and 25 after administration
- Concentration measurements using in-house ELISA techniques
  - RTX concentrations at each sampling time
  - ADA concentrations at day 25
- Pharmacokinetic modeling: two-compartment model, RTX–ADA interactions with irreversible binding (IB) model
- ADA concentrations, body weight (WT) and line tested as covariates




Figure 1: PK-ADA model



**Legends.** The kinetics of ADA input was described using 4 transit compartments, and zero and first-order rate constants for ADA production and transit, respectively. The kinetics of ADA was then tested as a two-compartment irreversible binding model, assuming that both rituximab and ADA share the same values for central volume of distribution, and transfer and elimination rate constants. Rituximab ADA-mediated elimination was assumed as occurring in both central and peripheral compartments.

**Parameters.**  $V_1$  central volume of distribution,  $k_{12}$  first-order central-to-peripheral ( $\approx 2.3$ ) and  $k_{21}$  peripheral-to-central transfer rate constants,  $k_{in}$  zero-order ADA input rate constant  $k_{tr}$  first-order transit and elimination rate constant and  $F_U$  antibody fraction unbound to neonatal Fc Receptor (FcRn) and eliminated.

- Modeling was carried out using nonlinear mixed-effect modeling software **Monolix Suite 2019** and **Simulx** (Lixoft®, Antony, France)

## CONCLUSION

- Nonlinear elimination may be imputable to immunization against rituximab only
  - Interactions of mAbs and ADA may be
    - similar to mAb-target interactions
    - described using TMDD modeling strategies
- Main issues may be the description of the onset of ADA production, hardly predictable in human therapy

## RESULTS

- Satisfactory description of concentration-time data with 2-compartment IB model
- Nonlinear elimination shape of concentrations observed in 2/3 mice
- Strong association between increasing ADA concentrations and increased  $k_{in}$  ( $\Delta AIC = -114.79$ )
- Mice lines presented different PK ( $V_1$ ,  $F_U$ ) and ADA production ( $k_{in}$ ) parameter values
- Onset of ADA-mediated elimination starting 7-10 days after rituximab injection

Table 1 : pharmacokinetic parameter estimates

| Parameter         | Unit              | Base model |      | Final model |      |
|-------------------|-------------------|------------|------|-------------|------|
|                   |                   | Estimate   | RSE% | Estimate    | RSE% |
| $V_1$             | mL                | 1.8        | 2.8  | 1.7         | 2.9  |
| Line(1)_ $V_1$    | —                 | —          | —    | 0.18        | 26   |
| Line(2)_ $V_1$    | —                 | —          | —    | -0.17       | 39   |
| BW_ $V_1$         | —                 | —          | —    | 0.34        | 30   |
| $F_U$             | —                 | 0.09       | 5.5  | 0.097       | 4.1  |
| Line(1)_ $F_U$    | —                 | —          | —    | -0.33       | 11   |
| Line(2)_ $F_U$    | —                 | —          | —    | 0.46        | 13   |
| $k_{12}$          | day <sup>-1</sup> | 2.2        | 4.7  | 2.3         | 3.7  |
| $k_{21}$          | day <sup>-1</sup> | 1.1        | 4.8  | 1.2         | 1.3  |
| $k_{in}$          | day <sup>-1</sup> | 13.1       | 33   | 46.2        | 23   |
| Ln(ADA)_ $k_{in}$ | —                 | —          | —    | 1.1         | 7.6  |
| Line_ $k_{in}$    | —                 | —          | —    | 2.2         | 13   |
| $k_{tr}$          | day <sup>-1</sup> | 0.24       | 6.6  | 0.21        | 2.4  |
| $k_{deg}$         | day <sup>-1</sup> | 0.11       | 2.8  | 0.83        | 1.6  |
| $\omega_{V1}$     | —                 | 0.28       | 6.7  | 0.18        | 7.3  |
| $\omega_{Fu}$     | —                 | 0.19       | 11   | 0.092       | 16   |
| $\omega_{Kin}$    | —                 | 3.3        | 9.8  | 1.24        | 9.3  |
| $\sigma_{prop}$   | —                 | 0.14       | 3.5  | 0.14        | 3.4  |




Figure 2: Diagnostic plots of final model




Figure 3: Influence of ADA input on RTX pharmacokinetics

### Legends.

**Table1.**  $V_1$  central volume of distribution,  $k_{12}$  first-order central-to-peripheral ( $\approx 2.3$ ) and  $k_{21}$  peripheral-to-central transfer rate constants,  $k_{in}$  zero-order ADA input rate constant  $k_{tr}$  first-order transit and elimination rate constant and  $F_U$  antibody fraction unbound to neonatal Fc Receptor (FcRn) and eliminated, RSE relative standard ratio, ADA, anti-drug antibodies, BW body weight,  $\omega$  interindividual standard deviation,  $\sigma_{prop}$  proportional residual standard deviation.

**Figure 2.** Top: observed vs. model-fitted RTX concentrations values in linear (left) and logarithmic (right) scales. Middle: individual reweighted (IWRES, left) and normalized prediction distribution error (NPDE, right) vs. Gaussian law. Bottom: Visual predictive checks (VPC) for 25, 50 and 75<sup>th</sup> concentration percentiles.

## REFERENCES

- Bensalem A, Ternant D. Clin Pharmacokinet. 2020;59:857-874.
- Ng CM, Loyet KM, Iyer S, Fielder PJ, Deng R. Eur J Pharm Sci. 2014;51:51-8.
- Liao KH, Udata C, Yin D, Sewell KL, Kantaridis C, Alvarez DF, Meng X. Br J Clin Pharmacol. 2020;86:2182-2191.
- Gibiansky L, Gibiansky E. Expert Opin Drug Metab Toxicol. 2009;5:803-12.
- Dua P, Hawkins E, van der Graaf PH. CPT Pharmacometrics Syst Pharmacol. 2015;4:324-37.