

Background and Objectives

Para-aminosalicylic acid (PAS) is a **WHO-recommended drug** (Group C) for rifampicin-resistant tuberculosis (RR-TB) treatment. Prevention of resistance in companion RR-TB drugs by PAS is dosage related and at **higher concentrations**, PAS may exhibit significant **bactericidal activity** in addition to its bacteriostatic properties. PAS is mainly (~ 80%) renally eliminated.

The **optimal dosing of PAS in children is uncertain.**

We describe the pharmacokinetics of the widely available **slow-release PAS formulation** in children treated for RR-TB, to identify any important covariates and to identify optimal drug dosing in children through simulations.

Methods

Data

- 27 children from South-Africa on at least 2 weeks of a PAS-containing individualized RR-TB regimen, dosed at 150-200 mg/kg in two divided doses.
- PAS slow-release granule formulation by PASER was used.
- Blood samples taken pre-dose and at 1, 2, 4, 8 and 11 h post PAS dose.

Assays

- Liquid-liquid extraction method using ethyl acetate, followed by LC-MS/MS detection. LLOQ was 0.391 mg/L.

Model

- NONMEM 7.4.4 with FOCE-I was used to develop the population PK model.
- PsN, Pirana and Xpose were used in the model building process for data exploration, visualization and creation of diagnostics.
- Allometric scaling was used to account for the effect of body size (using either total body weight or fat-free mass) and age was tested using a sigmoid Emax maturation model(1).
- Model-based simulations evaluated current 2019 WHO weight-band doses (200-300 mg/kg/day given as a once- or twice-daily dose) against the adult pharmacokinetic target of 50-100 mg/L for peak concentrations.

Results

Model

- PAS followed one-compartment kinetics with first-order elimination and transit-compartment absorption.
- The model detected a large inter-occasion variability (IOV) in bioavailability.
- Allometric scaling using body weight was included as a standard for paediatric populations (1), although the model showed no significant improvement after its inclusion (same using fat-free mass).
- Maturation parameters could not be robustly identified, as of the 10 (37%) children below 2 years of age, only 2 were below 1 year of age. Bayesian priors based on previously reported renal function (2), with 20% uncertainty, were therefore included in the model to improve simulation estimates in younger children.
- Increased PAS clearance was observed in both pharmacokinetic profiles from the only patient receiving efavirenz; this was retained in the model, as this interaction was previously reported in adults(3).
- No effect of renal function, sex, ethnicity, nutritional status, HIV status, other antiretrovirals (lamivudine, abacavir, lopinavir/ritonavir) or other RR-TB drugs was detected.

Table 2: Parameter estimates of the final model

Parameters	Typical value (95% CI)	BSV*/BOV** % (95% CI)
CL (L/h) <i>a</i>	9.15 (6.97, 11.7)	34.2* (21.8, 46.8)
Effect of efavirenz on CL in a single child (%)	+162 (+32, +351)	
V _c (L) <i>a</i>	38.7 (27.9, 53.5)	
MTT _b	4.41 (3.81, 4.92)	34.2** (26.4, 45.6)
NN () <i>b</i>	16.3 (10.2, 25.3)	
F ()	1- fixed	65.1** (48.2, 86.2)
Additive error (mg/L) <i>c</i>	0.0782 - fixed	
Proportional error (%)	34.2 (28.7, 41.1)	

a Reported for a child with a median weight of 13.3 kg. *b* Bayesian prior (ref 2).
c Fixed to improve model stability.




Figure 1. Visual predictive check. Lines represent the 5th, 50th, and 95th percentiles of the observed data, shaded areas are the model-predicted 95% confidence intervals for the same percentiles. Observed data are displayed as blue circles.

Table 1: Patient characteristics, n (%) or median) (IQR)

Characteristics	Children (n=27)
Age	3.87 (1.78-5.23)
Males	16 (59.3)
Weight (kg)	13.3 (10.9-17.0)
Black race	14 (51.9)
Z-score < -2 σ	
Weight-for-age	5 (18.5)
Length-for-age	10 (37.0)
No (%) HIV-positive	4 (14.8)
Creatinine (μ mol/L)	26 (20.5-31.5)
Antiretroviral treatment (n=4)	
LPV/RTV-based regimen	3 (75)
EFV-based regimen	1 (25)
Antituberculosis treatment	
Terizidone	27 (100)
Pyrazinamide	25 (92.6)
High-dose isoniazid	25 (92.6)
Ethambutol	21 (77.8)
Ethionamide	21 (77.8)
Amikacin	15 (55.6)
Clofazimine	11 (40.7)
Levofloxacin	10 (37.0)
Linezolid	10 (37.0)
Moxifloxacin	9 (33.3)
Capreomycin	6 (22.2)
Ofloxacin	3 (11.1)
Amoxicillin-clavulanic acid	2 (7.41)
Delamanid	1 (3.70)

a Weight- and height-for-age Z-scores of <-2 include underweight/stunted to severely underweight/stunted participants.

Simulations

- Target peak concentrations (50-100 mg/L) were achieved **only** in children receiving PAS at the **higher WHO dose** range of 300 mg/kg in a **once-daily** dose.
- Children should be dosed at the **higher range** of current WHO-recommended PAS doses and in a **once-daily** dose to optimize treatment.

PAS WHO Weight-based daily dosing (2019)




Figure 2. Simulated peak (Cmax) and minimum (Cmin) concentrations of PAS vs. weight in children receiving either the current WHO-recommended dose of 200mg/kg/day (upper panel), or 300 mg/kg/day (lower panel) given as either a once- (left panel) or twice-daily (right panel) dose. The dark-grey bands represents the desired (50–100 mg/L), and the minimum (1 mg/L) adult target ($\pm 10\%$).

Conclusions

- A transit compartment adequately describes absorption for the slow-release PAS formulation.
- **A once-daily dose of PAS at the higher end of the current WHO dose range of 300 mg/kg/day is recommended in children treated for RR-TB.**

References

1. Holford N, Heo Y, Anderson B. 2013. A pharmacokinetic standard for babies and adults. *J Pharm Sci* 102:2941–2952.
2. Rhodin MN, Anderson BJ, Peters AM, Coulthard MG, Wilkins B, Cole M, Chatelut E, Grubb A, Veal GJ, Keir MJ, Holford NHG. 2009. Human renal function maturation: A quantitative description using weight and postmenstrual age. *Pediatr Nephrol* 24:67–76.
3. De Kock L, Sy SKBB, Rosenkranz B, Diacon AH, Prescott K, Hernandez KR, Yu M, Derendorf H, Donald PR. 2014. Pharmacokinetics of para-aminosalicylic acid in HIV-uninfected and HIV-coinfected tuberculosis patients receiving antiretroviral therapy, managed for multidrug-resistant and extensively drug-resistant tuberculosis. *Antimicrob Agents Chemother* 58:6242–6250.

Acknowledgements:
The authors thank the children and parents for participating in this study. A special recognition to the University of Cape Town ICTS High Performance Computing team (<http://hpc.uct.ac.za>) for the use of their facilities.

Funding:
NIH R01 grant [Eunice Kennedy Shriver National Institute of Child Health and Human Development (NICHD-069169-01; PI: A. Hesselung), South African National Research Foundation SArCHI Chair (A. Hesselung), National Research Foundation, grant for rated researchers 109056 (P. Denti), H. Møllerion is supported by the Wellcome Trust (206379/Z/17/Z). UCT pharmacokinetic laboratory: drug assays were supported in part by the NIAID (UM1 AIO68634, UM1 AIO68636 and UM1A10106701, U01 AIO68632), the NICHD, and the National Institute of Mental Health (U01A068322). L. van der Laan was supported by the Swedish Foundation for International Cooperation in Research and Higher Education (STINT) jointly with the South African National Research Council, National Research Foundation [NRF] (NRF grant number: 101575).